Original Research Article

Fungal strains for mycological production of citric acid

Subedar Yadav^{1,*}¹Dept. of Chemistry, Sawatantrata Sangram Senani Vishram Singh Govt Post Graduate College, Chunar, Uttar Pradesh, India

ARTICLE INFO

Article history:

Received 12-05-2021

Accepted 19-05-2021

Available online 26-07-2021

Keywords:

Citric Acid

Aspergillus niger

NCIM

ABSTRACT

Citric acid has become an important raw material for general industrial use with many varied and expanding applications such as iron, steel, treatment and conditioning of industrial water supplies, preparation of alkyl resins, paints and in the printing of calico and textile industries. Mycological Production of citric acid includes preparation and sterilization of different media, culture medium, and seedling of culture tube, incubation of culture tubes, determination of citric acid formed and molasses left unfermented during the course of present investigation. The Present work deals mainly selection of potent strains of fungus.

© This is an open access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

1. Introduction

The production of citric acid by fungal strains is influenced by the carbohydrate sources such as molasses, concentration of molasses, incorporation of trace metals and some other compounds, monitoring of parameters to optimize the levels such as hydrogen ion concentration, temperature and incubation period. A probe is necessary for the above mentioned problems which will give the desired improved yield of citric acid.

Successful production of citric acid using potent strain of fungus is dependent on several factors of which the fungus, the culture conditions¹ and the raw materials used are most important. The success of a fermentation process depends to a very extent on the use of right type of fungal strain that can produce the desired end product at a minimum cost and in large quantities. The type of fungal strain used will depend on the product. *Aspergillus oryzae*² is a filamentous fungus used to saccharify rice, other grains and potatoes in making of alcoholic beverages. *Aspergillus flavus* is a fungal pathogen which causes post-harvest disease in cereal grains and legumes. *Aspergillus wentii* is used to process soybean. *Aspergillus foetidus* is used to produce koji for shochu- a

distilled Japanese alcoholic beverage and it is also utilized for the production of many useful enzymes that serve differing purposes. *Aspergillus niger*³⁻⁵ is a fungus and one of the most common species of the genus *Aspergillus*. It causes a disease called black mold on certain fruits and vegetables such as grapes, onions, peanuts and is a common contaminant of food.

Fungal strains of the *Aspergillus niger* group of molds have usually given most successful results both in the laboratory and on an industrial basis. Many of these molds produce high yields, imposes fairly uniform biochemical characteristics and they are easily cultivated and produce a negligible quantity of undesirable end products. In the present investigation the following potent fungal strains of some molds have been employed by the author for the mycological production of citric acid using Molasses as feedstock.

Table 1: Fungal strains employed for mycological production of citric acid

1(K)	<i>Aspergillus oryzae</i>	NCIM-647
2 (L)	<i>Aspergillus flavus</i>	NCIM-650
3 (M)	<i>Aspergillus wentii</i>	NCIM-661
4 (N)	<i>Aspergillus foetidus</i>	NCIM-511
5 (O)	<i>Aspergillus niger</i>	NCIM-683

* Corresponding author.

E-mail address: Subedar123@yahoo.co.in (S. Yadav).

NCIM stands for national Collection of industrial Microbes. The above citric acid producing strains of molds were employed for mycological production of citric acid and the observation are tabulated as in the Table 1 under heading result and discussion.

2. Experimental Method

The experiment was carried out with the above species of *Aspergillus* molds. Culture medium was prepared⁶ as follows:

Table 2:

Sucrose:	0.100g
Malt-Extract:	0.075g
Yeast extract:	0.075g
Peptone :	0.075g
Agar-Agar:	1.00g
Distilled Water:	0.100 ml
pH:	1.8

The pH of the culture medium was adjusted to 1.8 by adding requisite amount of KCl-HCl buffer solution. Malt extract is a sweet substances used as a dietary supplement⁷ for bacterial culture medium. Peptone mainly acts as a source of nitrogen and also carbon up to some extent. Agar-Agar is a gelatinous substances⁸ serves as the primary structural support for the algae's cell wall.

Microorganism grow in a vessel known as culture tubes. About a dozen of clean and steam sterilized dry culture tubes were filled 5 ml of culture medium each and kept in slanting position. A small quantity of *Aspergillus* was transferred to the freshly prepared culture tubes then kept in an autoclave maintained at 30 °C^{8,9} for 36 hrs. And then placed in refrigerator.

A small quantum of *Aspergillus* form culture tube was transferred to the freshly prepared inoculum medium^{10–16} in a conical flask and was kept in an incubator for about 36 hours for proper growth of fungus which is finally employed for the inoculation of production medium to get the citric acid on large scale. The composition of the production medium¹⁷ for the mycological production of citric acid by *Aspergillus* is as follows

Table 3:

Molasses:	20%
NH ₄ NO ₃ :	0.60%
KH ₂ PO ₄ :	0.60%
MgSO ₄ .7H ₂ O:	0.60%
pH:	1.8

After inoculation the fermentor flasks were placed in an incubator maintained at a constant temperature at 30 °C and were analyzed after 6, 8 and 10 days of incubation period for the production of citric acid and molasses unfermented.

When the citric acid fermentation is over the fermentation broth is kept at 50 °C to avoid contamination. The broth is drained off (filtered)¹⁸ and the mycelium mat is pressed to remove any acid contained in it. The amount of citric acid^{19–27} in the filtrate was measured by titration with 0.1M NaOH against phenolphthalein as an indicator.^{28,29} Total carbohydrates in the fermentation filtrate at various times was determined as glucose by anthrone-sulphuric acid method.^{30,31}

3. Result and Discussion

Table 4:

Sl. No	Isolates	Yield of citric acid * in g/100 ml
1(K)	<i>Aspergillus oryzae</i> NCIM-647	5.764
2(L)	<i>Aspergillus flavus</i> NCIM-650	5.983
3(M)	<i>Aspergillus wentii</i> NCIM-661	4.985
4(N)	<i>Aspergillus foetidus</i> NCIM-511	5.236
5(O)	<i>Aspergillus niger</i> NCIM-683	6.886

* Each value represents mean of three trials.

The data recorded indicates that the fungal strain designated (O) ie *Aspergillus niger* NCIM-683 has significant yield of citric acid i.e. 6.886 g/100ml in comparison to other isolates. It is thus concluded that all the citric acid producing fungus are not of equal importance though they are morphologically similar to each other. The author therefore has selected *Aspergillus niger* NCIM-683 for mycological production of citric acid for further study along with different mutagen.^{32–36}

4. Source of Funding

None.

5. Conflict of Interest

None.

6. Acknowledgement

I express my sincere gratitude to Professor SP Singh Department of Chemistry Magadh University Bodh Gaya for all support and guidance.

References

- William AR, Weber DJ. CDC-Guideline for Disinfection and Sterilization in Healthcare Facilities. Prions; 2010. Available from: <https://www.cdc.gov/infectioncontrol/pdf/guidelines/disinfection-guidelines-H.pdf>.
- Rokas A. The effect of domestication on the fungal proteome. *Trends Genet.* 2009;25(2):60–3. doi:10.1016/j.tig.2008.11.003.
- Samson RA, Houbaken J, Summerbell RC, Flannigan B, Mille JD. Common and important species of fungi and

- actinomycetes in indoor environments; 2016. Available from: <https://pure.knaw.nl/portal/en/publications/common-and-important-species-of-fungi-and-actinomycetes-in-indoor>.
4. Abarca ML, Bragulat MR, Castellá G, Cabañes FJ. Ochratoxin A production by strains of *Aspergillus niger* var. *niger*. *Appl Environ Microbiol*. 1994;60(7):2650–2. doi:10.1128/aem.60.7.2650-2652.1994.
 5. Schuster E, Coleman ND, Frisvad J, Dijck PV. On the safety of *Aspergillus niger* - a review. *Appl Microbiol Biotechnol*. 2002;59(4):426–35. doi:10.1007/s00253-002-1032-6.
 6. Ratnasamy P. Maintenance of Culture 2nd Edition NCL Pune. India; 1977.
 7. British pharmaceutical codex. Pharmaceutical Society of Great Britain; 1907. Available from: <https://wellcomecollection.org/works/mfze4eb8>.
 8. Vandenberghe LPS, Soccol CR, Pandey A, Lebeault JM. Microbial production of citric acid. *Braz Arch Biol Technol*. 1999;42(3):263–76. doi:10.1590/s1516-89131999000300001.
 9. Mohsen Y, Agarwal M. Effect of *Celastrus paniculatus* on Learning, Memory and Serum Biochemistry of Aging Albino Rats. *Int J Chem Chem Eng*. 2013;3:181–90.
 10. Zimmermann H, Lindgren A, Schuhmann W, Gorton L. Anisotropic Orientation of Horseradish Peroxidase by Reconstitution on a Thiol-Modified Gold Electrode. *Chem Eur J*. 1993;95:322–49.
 11. Koemann RH. I.A Modification Of The Miller-Muntz Method For Colorimetric Determination Of Lactic Acid. *J Biol Chem*. 1940;135(1):105–9. doi:10.1016/s0021-9258(18)73164-5.
 12. Rusdell LJ. The Nature of Metaphysical Thinking. *Nature*. 1945;156(3956):217–8. doi:10.1038/156217a0.
 13. Villano ZF, Langella G. [Determination of glycoproteins in blood in man by means of paper electrophoresis. III. Results in pathological cases and comparison with blood protein changes]. *Biol Soc Ital Biol Sper*. 1943;32(6):349–51.
 14. Horold CE. Studies in the Genus *Eremascus*: I. The Rediscovery of *Eremascus albus* Eidam and Some New Observations concerning its Life-history and Cytology. *Ann Bot*. 1950;14:127–48.
 15. Turyanitsa ID, Oleksyuk ID, Kozmanko II. ChemInform Abstract: SYSTEM SB2TE3-SBJ3 UND EIGENSCHAFTEN DER VERBINDUNG SBTEJ. *Chemischer Informationsdienst*. 1973;11(4):4. doi:10.1002/chin.197346018.
 16. Garrett SD. Biology of Root infecting fungi; 1956. p. 292. Available from: https://journals.iwv.com/soilsci/Citation/1956/07000/Biology_of_Root_Infecting_Fungi.11.aspx.
 17. Currie JN. The Citric Acid Fermentation Of *Aspergillus Niger*. *J Biol Chem*. 1917;31(1):15–37. doi:10.1016/s0021-9258(18)86708-4.
 18. Yokoya F. Citric Acid Production. *Ind Fermentation Ser*. 1992;4(1):1–82.
 19. Grewal HS, Kalra KL. Fungal production of citric acid. *Biotechnol Adv*. 1995;13(2):209–34. doi:10.1016/0734-9750(95)00002-8.
 20. Pandey A, Soccol CR, Leon JA, Nigam P. Production of Organic Acids by Solid-State Fermentation; 2001. p. 113–26. Available from: <https://pure.ulster.ac.uk/en/publications/solid-state-fermentation-in-biotechnology-fundamentals-and-applic-2>.
 21. Vandenberghe LPS, Soccol CR, Pandey A, Lebeault JM. Review: Microbial production of citric acid. *Braz Arch Biol Technol*. 1999;p. 42–263.
 22. Yokoya F. Citric Acid Production. In: Industrial Fermentation Series. *Int J Pharm Pharm Sci*. 1992;7:1–82.
 23. Kubicek CP, Röhr M, Rehm HJ. Citric Acid Fermentation. *Crit Rev Biotechnol*. 1985;3(4):331–73. doi:10.3109/07388558509150788.
 24. Jianlong W, Xianghua W, Ding Z. Production of citric acid from molasses integrated with in-situ product separation by ion-exchange resin adsorption. *Bioresource Technol*. 2000;75(3):231–4. doi:10.1016/s0960-8524(00)00067-5.
 25. Friesen DT, Babcock WC, Brose DJ, Chambers AR. Recovery of citric acid from fermentation beer using supported-liquid membranes. *J Membrane Sci*. 1991;56(2):127–41. doi:10.1016/s0376-7388(00)80803-0.
 26. Roukas T, Kotzekidou P. Pretreatment of date syrup to increase citric acid production. *Enzyme Microbial Technol*. 1997;21(4):273–6. doi:10.1016/s0141-0229(97)00041-0.
 27. Legisa M, Gradisnik-Grapulin M. Sudden substrate dilution induces a higher rate of citric acid production by *Aspergillus niger*. *Appl Environ Microbiol*. 1995;61(7):2732–7. doi:10.1128/aem.61.7.2732-2737.1995.
 28. Suzuki A, Sarngbin S, Krimura K, Usami S. Direct production of citric acid from starch by a 2-deoxyglucose-resistant mutant strain of *Aspergillus niger*. *J Ferment Bioeng*. 1996;81(4):320–3. doi:10.1016/0922-338X(96)80584-8.
 29. Sakurai A, Imai H, Ejiri T, Endoh K, Usami S. Citric acid production by surface culture using *Aspergillus niger*: Kinetics and simulation. *J Fermentation Bioeng*. 1991;72(1):15–9. doi:10.1016/0922-338X(91)90139-8.
 30. Hamamci H, Hang YD. Production of citric acid by immobilized dried reactivated *Aspergillus niger*. *Biotechnol Tech*. 1989;3:51–4. doi:10.1007/bf01876221.
 31. Tan KH, Ferguson LB, Carlton C. Conversion of cassava starch to biomass, carbohydrates, and acids by *Aspergillus niger*. *J Appl Biochem*. 1984;6(1):80–90.
 32. Yadav S. Mycological Production of citric acid by *Aspergillus niger* exposed to 3-(α -acetylbenzyl)-4-hydroxycoumarin sodium salt. *J Environ Sci Comput Sci Eng Technol*. 2018;7(1):322–8. doi:10.24214/jecet.A.7.1.05459..
 33. Yadav S. Mycological Production of Citric Acid Exposed to 3-Acetamidocoumarin. *J Chem Pharm Res*. 2017;7(1):54–9.
 34. Rokas A. The effect of domestication on the fungal proteome. *Trends in genetics: TIG 2009*;p. 60–63.
 35. Samson RA, Houbraken J, Flannigan B, Miller JD. Common and important species of fungi and actinomycetes in indoor environments in Microorganisms in Home and Indoor Work Environments; 2001. p. 287–92.
 36. Cyclopedia of India and of eastern and southern Asia, commercial . vol. 2; 1871. p. 83. Available from: <https://www.biodiversitylibrary.org/item/211446#page/4/mode/1up>.

Author biography

Subedar Yadav, Senior Assistant Professor

Cite this article: Yadav S. Fungal strains for mycological production of citric acid. *Int J Pharm Chem Anal* 2021;8(2):59-61.